



TOPICAL DILTIAZEM FOR PAIN RELIEF AFTER HEMORRHOIDECTOMY

Dr Sandeep Rao	M.S., Associate Professor Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Mohan Lal Didel	M.S. Senior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Adeesh Jain	M.S., Additional Professor, Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Milind Patil	M.S., Assistant Professor, Department of General Surgery, S.S.G.Hospital, Vadodara.
Dr Ashvin Kankotiya	M.S., Assistant Professor, Department of General Surgery, S.S.G.Hospital, Vadodara.
Dr Rahul Rathwa	M.S. Senior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Anand Ratra	M.B.B.S. ,Third year Junior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Rahul Masne	M.B.B.S. ,Third year Junior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.
Dr Kirtan Gosai	M.B.B.S. ,Third year Junior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.

ABSTRACT

Aim: To assess effectiveness of 2% Topical Diltiazem ointment use for pain relief after open haemorrhoidectomy surgery.

Methods: A randomized, prospective, randomised study was conducted on 2 groups of patients in the postoperative period of open haemorrhoidectomy. Each group consisted of 46 patients. Group A received topical diltiazem in the anal region 3 times a day and group B received a tablet tramadol.

Results: In group A, the mean score on the visual analogue scale was 5.6 at 24 h, 1.5 at 48 h, and 0.5 at 72 h. In group B, it was 6 at 24 h, 3.6 at 48 h, and 2 at 72 h

Conclusion: In this study average pain scores after using 2% Topical Diltiazem ointment at 48 hours was 1.5 as compared control group 3.6 and at 72 hours was 0.5 as compared control group 2.0 which is nearly comparable to above mentioned studies. hence, there was significant reduction of haemorrhoidectomy pain at 48 hours and 72 hours in patients who applied 2% Topical Diltiazem ointment.

KEYWORD

open haemorrhoidectomy, topical diltiazem, post-operative pain

*Corresponding Author Dr Anand Ratra

M.B.B.S. ,Third year Junior Resident, Department of General Surgery, S.S.G. Hospital, Vadodara.
Email id- anandratra1234@gmail.com

INTRODUCTION

Haemorrhoids are cushions of submucosal tissue containing venules, arterioles, and smooth muscle fibers that are located in the anal canal. There are few diseases more chronicled in human history than symptomatic haemorrhoidal disease. References occur in ancient texts dating back to Babylonian, Egyptian, Greek, and Hebrew cultures.^{1,2} Included in many of these writings are multiple recommended treatment regimens, including anal dilation, topical ointments, and the intimidating red hot poker.^{3,4} Although few people have died of hemorrhoidal disease, many patients wish they had, particularly after therapy, and this fact led to the beatification of St. Fiachre, the patron saint of gardeners and haemorrhoidal sufferers.⁵ It is important for the practitioner to give a more humane approach to hemorrhoidal disease, with the emphasis on cost-effectiveness with minimal morbidity and mortality.

Because haemorrhoids are a normal part of anorectal anatomy, treatment is only indicated if they become symptomatic. Today there are a variety of treatments available for symptomatic hemorrhoids. Patients can be treated conservatively with seitz bath, anaesthetics, corticosteroids, analgesics, soothing agents.

Office procedures for treatment include: rubber band ligation, injection therapy, cryotherapy, and photocoagulation.

Surgical methods like Open Haemorrhoidectomy, Whitehead's Haemorrhoidectomy, Procedure for Prolapse Haemorrhoids/Stapled Haemorrhoidectomy⁶ and Doppler-Guided Haemorrhoidal Artery Ligation⁷ have evolved over the period of years.

Spasm of the internal anal sphincter is thought to be related to the source of pain after haemorrhoidectomy and may be a component of postoperative haemorrhoidectomy pain.

Diltiazem, a calcium channel blocker, has been shown to reduce resting anal pressure and relax gastrointestinal smooth muscle⁽⁸⁻¹¹⁾

Topical application of diltiazem is commonly used to relieve pain after haemorrhoidectomy.

OBJECTIVES OF THE STUDY

AIM:

- To assess effectiveness of 2% Topical Diltiazem ointment use for pain relief after open haemorrhoidectomy surgery.

INCLUSION CRITERIA:

1. All patients posted for Open Milligan-Morgan haemorrhoidectomy.

EXCLUSION CRITERIA:

- Patient already under treatment with calcium channel blocker agent
- Patient that did not give their informed consent to get involved in this study
- Patient age \leq 18 yrs
- Patient with
 - Severe high blood pressure
 - Orthostatic hypotension

MATERIALS AND METHODS

The study has been done in Dept. of Surgery, Medical College Baroda and SSG Hospital, the period of which has extended from March 2018 to Dec 2019

Duration of Study:

One year 10 months, (March 2018 to Dec 2019)

Type of Study: Prospective Randomized Control trial

Sample Size: A total of 92 patients were taken in this study out of which 46 were allocated to the case group and 46 to the control group.

METHODS

All patients undergoing Open Milligan Morgan haemorrhoidectomy was divided in two group

Group A(ODD) -was given Topical Diltiazem 2% and rescue analgesic for pain relief.

Group B(EVEN)-was given analgesic, Tablet Tramadol 50 mg 1BD for 3 days.

1gm Diltiazem ointment 2% was measured by spoon and applied to the patient by surgeon's index finger and applied circumferentially over haemorrhoidectomy wound, immediate post-operative time in OT before shifting patient to recovery room. Then next day morning in ward Diltiazem ointment 2% was applied over haemorrhoidectomy wound by resident doctor thrice a day in left lateral position for 3 days.

Pain intensity was measured by visual analogue scale (VAS) as shown in figure 1.

Pain assessment was done at 24hrs, 48hrs and 72 hrs after the end of the surgical procedure.

Rescue medication: tablet Tramadol 50 mg 1 SOS

Indication of rescue medication: VISUAL ANALOGUE SCALE score \geq 6.

Patient was instructed to remain in supine position for

some time and watch for any complication if occurs then report to the doctor.

RESULTS

Total number of 92 patients were taken into the study and were then divided into cases and controls. 46 patients were taken into each group.

AGE DISTRIBUTION

It was observed that majority of the patients (30.43%) were in the 31-40 years of age group. Distribution of patient according to the age is as shown in [Table 1].

This suggests that in this age group, patients were motivated to undergo operative procedure for symptomatic hemorrhoids.

Relation of Age to VAS scores in Case group:

It was observed that 21-30 years and 61-70 years age group had slightly higher VAS scores as compared to other age groups as shown in table 2.

Relation of Age to VAS scores in Control group:

It was observed that 21-40 years age group had higher VAS scores as compared to other groups as shown in table 3.

VAS SCORE ANALYSIS

At 24 hours intervals there was no significant difference seen. Only 9 patients had VAS score more than 6 and they needed rescue medicine (Tablet Tramadol 50mg) only one time.

At 48 hours and 72 hours, VAS Scores were checked and it was found that the control group clearly had a higher number. This was a significant difference as shown in table 4.

It is therefore clearly observed that the patients who were applied 2% Topical Diltiazem experienced little or no pain. However, on the other hand the control group in which regular analgesic was given, complained of post-operative pain.

Inference:

At 24 hours intervals there was no significant difference seen. At 48 hour and 72 hours VAS, the p value was calculated to be less than 0.05 which shows a significant difference between the case and control groups. The patients applied 2% Topical Diltiazem ointment had lower pain scores as compared to those who received Tablet Tramadol as shown in table 5.

Other Complications

It is seen that urinary retention was seen more in patients the control group as compared to the cases group.

This can be attributed to the fact that pain led to urinary retention in the control group. Two of these patients required catheterization. However, the catheter was removed the following day with spontaneous passing of urine.

It is therefore advisory to have a good post-operative analgesia for patients undergoing surgery for hemorrhoids.

Bleeding was seen in one patient in the control group. It was only a few milliliters and did not require transfusion. Bleeding stopped with adequate compression followed by padded dressing at bedside.

Post-operative nausea and vomiting was seen in three patients in the control group and none in the cases group.

It was controlled on giving antiemetic (Tablet Domperidone 10 mg sos).

Light headache was seen in 2 patients and dizziness was seen in 2 patient of case group.

DISCUSSION

Hemorrhoidectomy is usually associated with considerable pain during the postoperative period, which may delay discharge, recovery, and return to work. Accordingly, a variety of methods for reducing pain after hemorrhoidectomy have been investigated, including pudendal nerve block, perianal infiltration of local anesthetics or botulinum toxin, and application of topical preparation such as metronidazole, glyceryl nitrate, and calcium channel blocker.²²

Diltiazem, a hydrophilic benzothiazepine, is a L-type calcium channel antagonist given orally for treatment of various cardiovascular disease.²³

Postoperative pain and defecation difficulty immediately after surgery are major clinical problems following hemorrhoidectomy.

The meta-analysis demonstrated that the use of Topical Diltiazem was associated with a more significantly improvement and relief of anal pain after hemorrhoidectomy and a lower use of additional oral pain killers with fewer side effects.

Higher resting anal pressure was believed to be present in patient with anal fissure and haemorrhoids²⁴⁻²⁵ painful spasm of internal anal sphincter leading to high pressure zone in anal canal is also thought to be the source of pain after anal surgery and may contribute to post haemorrhoidectomy pain.²⁶

Maintenance of smooth muscle tone of internal sphincter is a calcium dependent mechanism and the use of calcium channel blockers causes relaxation of gastrointestinal smooth muscle.⁸⁻¹¹

Diltiazem acts by inhibiting the flow of extracellular calcium ions into the sarcoplasm of internal anal sphincter with a consequent saving of oxygen, resulting in muscle relaxation and pain relief.²⁷ it would therefore seem reasonable to apply medication aimed at reducing internal sphincter pressure which might be beneficial in reducing anal pain after haemorrhoidectomy.

This metanalysis has shown that the post-operative use of 2% Diltiazem ointment is statistically significant in reducing post-operative pain at 48 hours and 72 hours.

This was confirmed by carapeti et al.who showed that a single dose of 60mg diltiazem lowered the maximum resting anal sphincter pressure by a mean of 21%.²⁰

Amoli and associates prospectively evaluated a role of topical 2% diltiazem application after haemorrhoidectomy and concluded that perianal application of diltiazem ointment after haemorrhoidectomy significantly reduces postoperative pain.¹⁸

Chauhan et al prospectively randomised 108 patients of Third- and fourth-degree haemorrhoids and assessed the efficacy of internal sphincterotomy compared with topical application of 2% Diltiazem ointment after haemorrhoidectomy for pain relief. They encountered no significant difference in mean pain scores on second postoperative day as both the procedure relieve the spasm of anal sphincter.²¹

Silverman et al evaluated the effect of Topical Diltiazem in reducing pain after haemorrhoidectomy. They randomly assigned 18 patients of haemorrhoidectomy to receive 2% Diltiazem and placebo ointment and noticed mean pain scores of 5.2 in diltiazem group as compared to 8.8 in placebo group on second postoperative day.¹⁹

In this study average pain scores after using 2% Topical

Diltiazem ointment at 48 hours was 1.5 as compared control group 3.6 and at 72 hours was 0.5 as compared control group 2.0 which is nearly comparable to above mentioned studies. hence, there was significant reduction of haemorrhoidectomy pain at 48 hours and 72 hours in patients who applied 2% Topical Diltiazem ointment.

The control group had more patients who had a complaint of post-operative nausea and vomiting due to side effect of Tablet Tramadol.

Some advantages of the current study are worth highlighting. We included only elective, **Open Milligan Morgan Hemorrhoidectomy** in our study to avoid the type, nature, and duration of pain associated with different types of surgery. Moreover, all the observations were carried out by a single observer to eliminate any inter observer variability. Thus, we can assume that the difference in pain relief reflects only the efficacy of the antinociceptive measures.

VAS Score Analysis

In the current study it is clearly seen that VAS scores are lower in the group which was given 2% Topical Diltiazem. Patients in the control group felt post-operative pain in the first 72 hour of post-operative time. This also led to the increased incidence of use of tablet tramadol in the control group.

Therefore, in line with the previous study, this study also had results which confirms the efficacy on Topical Diltiazem 2% in reducing the VAS scores.

Age Distribution

In this study, majority of the patients belonged to 31-40 years of age group.

It was observed that VAS scores were higher in age group of 21-30 years and 61-70 years in case group.

In the control group VAS scores were higher in 21-40 years age group.

In the study conducted by S Suchdev et al control group had mean age group of 39.85(±14.91) and case group had mean age of 37.73(±14.90); difference between two means was statistically insignificant (p-value=0.637)

There was a slight difference in the age group as compared to the previous studies showing predominance in the 31-50 years of age group.

Other postoperative complications

In this study case group had one patient with urinary retention for which he was catheterized. On the other-hand control group had 3 patients with urinary retention post operatively which he was catheterized.

Bleeding was seen in one patient of the control group and none in the case group.

Post-operative nausea and vomiting was seen in 3 patients of control group and none in the case group.

This shows that there was a higher rate of complications possibly pertaining to pain in the control group as compared to those in the case group.

ACKNOWLEDGEMENTS

No funding was received.

DECLARATIONS

Funding:None

Conflict of interest:

Nondeclared

Ethical approval:required and approved

SUMMARY

This study 92 patients were taken who were distributed equally into two groups.

From this study it is evident that the effect of 2% Topical Diltiazem ointment is a positive one and majority of the patients who were in this group didn't complain of pain post operatively. On the contrary, in the control group the patients invariably complained of pain and had to be given pain medications.

In this study at 24 hours intervals there was no significant difference seen.

At 48 hours and 72 hours, VAS scores were checked and it was found that the control group clearly had a higher number. This was a significant difference.

Majority of the patients were in the age group of 31-40. The patients who participated in this trial were mainly those who had grade 3 and grade 4 symptomatic hemorrhoids.

Patients in the control group seemed to have higher post-operative urinary retention and post-operative nausea and vomiting rates. This finding could be possibly attributed to pain.

The VAS scores observed clearly indicates the efficacy of 2% Topical Diltiazem in preventing post-operative pain in open hemorrhoidectomies.

TABLES

Table 1: Age wise distribution of patients

AGE (YEARS)	NO. OF PATIENTS	CASES	CONTROLS	PERCENTAGE
21-30 y	14	6	8	15.21%
31-40 y	28	18	10	30.43%
41-50 y	20	12	8	21.73%
51-60 y	18	6	12	19.56%
61-70 y	12	4	8	13.04%

Table 2: Relation of Age to VAS scores in Case group (average VAS score)

AGE (YEARS)	VAS @24HOUR	VAS @48HOUR	VAS @72HOUR
21-30 y	5.83	1.67	0.50
31-40 y	5.61	1.44	0.44
41-50 y	5.50	1.75	0.58
51-60 y	5.17	1.0	0.50
61-70 y	5.75	1.5	0.50

Table3: Relation of Age to VAS scores in Control group (average VAS score)

AGE (YEARS)	VAS @24HOUR	VAS @48HOUR	VAS @72HOUR
21-30	6.38	3.88	2.50
31-40	6.30	3.80	2.00
41-50	5.63	3.50	1.88
51-60	5.75	3.33	1.83
61-70	6.0	3.63	1.88

Table 4: Vas Scores(average)

	VAS at 24hour	VAS at 48hour	VAS at 72hour
Case	5.6	1.5	0.5
Control	6	3.6	2

Table 5: Analysis

	Group	N	Mean	SD	P-VALUE
VAS24	Case	46	5.56	1.13	0.085
	Control	46	6	1.26	
	Total	92			
VAS48	Case	46	1.5	1.03	<0.00001

	Control	46	3.6	0.99	
	Total	92			
VAS72	Case	46	0.5	0.51	<0.00001
	Control	46	2.00	0.84	
	Total	92			

FIGURES




Figure 1: pain assessment tool

CONCLUSION

The analgesic effect of local 2% Topical Diltiazem ointment contributes to improvement in pain control after open hemorrhoidectomy. The technique is simple and easy to learn. Further it can be performed in a few minutes. Diltiazem have fewer side effects profile making it a safe drug for this use. Hence, pain which is a major complication of

OPEN MILLIGAN MORGAN HAEMORRHOIDECTOMY surgeries can be easily dealt with this method of pain control.

We conclude by recommending that 2% topical diltiazem ointment is not only effective but is also a safe procedure, and can be a better alternative strategy for reducing the pain of patients who are undergoing **OPEN MILLIGAN MORGAN HAEMORRHOIDECTOMY**.

REFERENCES

- Holley CJ: History of hemorrhoidal surgery. South Med J 1946;39:536.
- Madoff RD: Biblical management of anorectal disease. Presented at the Midwest Society of Colon and Rectal Surgeons. March, 1991; Breckenridge, Colorado.
- Dirckx JH: The biblical plague of "hemorrhoids". An outbreak of bilharziasis. Am J Dermatopathol 1985;7:341.
- Maimonides M, Rosner F, Munter S [trans]: Treatise on hemorrhoids. Philadelphia, 1969, JB Lippincott.
- Rachochot JE, Petourand CH, Riovoire JO: Saint Fiacre: The healer of hemorrhoids and patron saint of proctology. Am J Proctol 1971;22:175.
- Manfredelli S, Montalto G, Leonetti G, et al. Conventional (CH) vs. stapled hemorrhoidectomy (SH) in surgical treatment of hemorrhoids. Ten years experience. Ann Ital Chir. 2012;83:129-134.
- Schuurman JP, Borel Rinkes IH, Go PM. Hemorrhoidal artery ligation procedure with or without Doppler transducer in grade II and III hemorrhoidal disease: a blinded randomized clinical trial. Ann Surg. 2012; 255:840-845.
- Carapeti EA, Kamm MA, Evans BK, Phillips RK. Topical diltiazem and bethanechol decrease anal sphincter pressure without side effects. Gut. 1999;45:719-722.
- Jonard P, Essami B. Diltiazem and internal anal sphincter. Lancet. 1987;1:754.
- Staneva-Stoytcheva D, Venkova K. Effect of calcium antagonists diltiazem, verapamil and nitrendipine on the contractile responses of guinea-pig isolated ileum to electrical stimulation or carbachol. J Pharm Pharmacol. 1992;44:321-325.
- Morales olivas FJ, Cortijo j, Esplugues jv, et al. Effect of verapamil and diltiazem on isolated gastro-oesophageal sphincter of the rat. J Pharm Pharmacol. 1985;37:208-209.
- Ellesmore S, Windsor AC. Surgical History of Haemorrhoids. In: Charles MV, editor. Surgical Treatment of Haemorrhoids. London: Springer; 2002; 1-4.
- Shampo MA. Brief history of colorectal surgery. J Pelvic Surg. 1998;4:259-267.
- Milligan ET, Morgan CN, Jones LE, Officer R. Surgical anatomy of the anal canal and operative treatment of haemorrhoids. Lancet. 1937;11:1119-94.
- Adotey JM, Jebbin NJ. Anorectal disorders requiring surgical treatment in the University Teaching Hospital, PortHarcourt. Niger J Med. 2004;13:350-4.
- Uba AF, Inezue CH, Obekpa PO, Iya D, Legbo JN. Open haemorrhoidectomy revisited. Niger J Med. 2001;10:185-8.
- Uba AF, Obekpa PO, Ardill W. Open versus closed haemorrhoidectomy. Niger Postgrad Med J. 2004;11:79-83.
- Amoli HA, Notash AY, Shahandasti FJ, et al. A randomized, prospective, double blind, placebo-controlled trial of the effect of topical diltiazem on post haemorrhoidectomy pain. Colorectal Dis. 2011 Mar; 13 (3):328-332
- Silverman R, Bendick PJ, Wasvary HU. A randomized, prospective, double-blind, placebo-controlled trial of the effect of a calcium channel blocker ointment on pain after hemorrhoidectomy. Dis Colon Rectum. 2005 Oct; 48

- (10):1913-1916.
20. Carapeti EA, Kamm MA, Evans BK, Phillips RK. Topical diltiazem and bethanechol decrease anal sphincter pressure without side effects. *Gut*. 1999;45:719-722.
 21. Chouhan A, Thomas S, Bishnoi PK, Hadke NS. Randomised controlled trial to assess the role of raised anal pressure in the pathogenesis of symptomatic early hemorrhoids. *Dig Surg*. 2007;24(1):28-32.
 22. Cheetham MJ, Phillips RK. Evidence-based practice in haemorrhoidectomy. *Color Dis*. 2001;3:126-134.
 23. Triggle DJ. Calcium channels, and calcium channel antagonist. *Can J Physiol Pharmacol*. 1990;68:1474-1481.
 24. Arabi Y, Alexander Williams J, Keighley MR. Anal pressure in haemorrhoids and anal fissure. *Am J Surg*. 1977;134:608-610.
 25. Patti R, Almasio PL, Arcara M, et al. Long-term manometric study of anal sphincter function after hemorrhoidectomy. *Int J Colorectal Dis*. 2007 mar; 22c(3):253-257.
 26. Loder PB, Kam J, Phillips RK. Haemorrhoids: pathology, pathophysiology and aetiology. *Br J Surg*. 1994;81:946-954.
 27. Cook TA, Brading AF, Mortensen NU. Differences in contractile properties of anorectal smooth muscle and the effect of calcium channel blockade. *Br J Surg*. 1999;86:70-75.
 28. Prasad ML, Abcarian H. Urinary retention following operations for benign anorectal diseases. *Dis Colon Rectum* 1978;21:490-2.
 29. Khubchandani IT. Internal sphincterotomy with hemorrhoidectomy does not relieve pain: A prospective, randomized study. *Dis Colon Rectum* 2002; 45: 1452-7